

# Effect of seeding rate on organic canola emergence, red legged earthmite (*Halotydeus destructor*) density, weed biomass and grain yield in northeast Victoria.

V.F. Burnett<sup>1</sup>, J.H. Schneider<sup>1</sup>, F.W. Dempsey<sup>1</sup> and B. Hemantha Rohitha<sup>2</sup>

<sup>1</sup> Department of Natural Resources and Environment, Rutherglen Research Institute, RMB 1145, Chiltern Valley Road, Rutherglen Victoria 3685

<sup>2</sup> Department of Natural Resources and Environment, Victorian Institute for Dryland Agriculture, Private Bag 260, Horsham, Victoria 3401

## Abstract

An experiment to investigate the effect of seeding rate on canola emergence, red legged earth-mite (RLEM) density, weed biomass and grain yield was established at Rutherglen Research Institute in 2001 under a certified organic regime. Canola, cv. Ripper, was sown in acid soil (pH<sub>Ca</sub> 5.2) at three seeding rates; 5, 7.5 and 10 kg/ha with phosphorus (P) applied with the seed as Guano™ at 10 kg/ha. Increasing seeding rate resulted in more canola plants/m<sup>2</sup>, less weed biomass and improved yields. Seeding rate did not affect the number of ryegrass seedlings at emergence or crop biomass. Red legged earth-mite numbers were highest on the 10 kg/ha seeding rate but there were no significant differences between the treatments. Increasing seeding rate had no effect on protein or oil contents. No effective management of the RLEM population was carried out during 2000 so the risk of the canola being affected by this pest was high. Despite higher RLEM densities on the 10 kg/ha seeding rate earlier in the season, the increased plant density was able to survive the mite attack and significantly reduce weed biomass to provide economic yields. Increased seeding rates in canola may provide an effective tool for organic producers to reduce the impact of weeds and RLEM.

## Keywords

Organic farming, canola, seeding rate, ryegrass, red legged earthmite, grain yield

## Introduction

Canola has been identified as having strong potential for export growth within the organic sector (1). Conventional canola production is heavily reliant on inputs of herbicides and insecticides to manage weeds and insect pests such as RLEM (*Halotydeus destructor*). These inputs are not allowed in a certified organic farming system, so an integrated approach to management of pests is required. Previous research by Peltzer (2) in Western Australia has shown canola to be a poor competitor against ryegrass in comparison to cereals and yield losses of between 9 and 30% have been shown with ryegrass densities of 300 plants/m<sup>2</sup> (3). This experiment was established to determine whether increasing the seeding rate of canola could affect competition from ryegrass, and withstand RLEM attack under a certified organic production system.

## Methods

The experiment was established at Rutherglen Research Institute in 2001. The site was a mix of Chromosol and Dermosol soils. Soil chemical properties measured in the 0-10 cm depth were pH<sub>Ca</sub> 5.2, 20 mg/kg of available sulphur, and 4 mg/kg Olsen P. Available nitrogen of 121.2 kg/ha was measured in the 0-100 cm depth. Canola, cv. Ripper, was sown into a prepared seed-bed at three rates; 5, 7.5 and 10 kg/ha with phosphorus applied with the seed as Guano™ at 10 kg/ha and the experiment was replicated 4 times. Difficulty in obtaining sufficient precision between the 5 and 7.5 kg/ha seeding rate was experienced with the machinery used in sowing. This has affected the results obtained for canola emergence at the low seeding rates such that comparisons can only be made between the two lower rates and the 10 kg/ha rate. Growing season rainfall (April - November) was 318 mm compared with a long-term average of 439 mm. Previous weed management on the site involved cutting a legume hay crop in October 2000. The canola was sown in early June, windrowed in November, and harvested in

early December. At crop emergence 10 samples per plot (28m<sup>2</sup>) were taken to assess RLEM population density using a suction sampler with a diameter of 125 cm in mid June 2001.

## Results

Increasing seeding rate to 10 kg/ha resulted in more canola plants/m<sup>2</sup> at emergence (Table 1). The pre-sowing ryegrass density on this site was 1,000 plants/m<sup>2</sup> in early April. Cultivation prior to seeding resulted in effective ryegrass control but an average of 91 ryegrass plants/m<sup>2</sup> germinated with the canola (Table 1), and this had increased to a density of 108 plants/m<sup>2</sup> by early flowering (Table 2). RLEM density on this site in Spring 2000 was 7,890 mites/m<sup>2</sup> and when we measured the density in June 2001, there was an average of 76 mites/m<sup>2</sup>. When canola and weed biomass were assessed in late October, significant differences were found in the amount of weed dry matter between the lower, and the high seeding rates (Table 2). The higher seeding rate resulted in a lower weed biomass measurement, and this trend was repeated in grain yield, with the 10 kg/ha rate resulting in the highest yield of 1.6 t/ha (Table 2). There was no effect of seeding rate on oil content or grain protein (Table 2). Ryegrass density throughout the season was consistently below 300 plants/m<sup>2</sup> where significant yield losses in canola can occur (3), and the density of RLEM was below control thresholds currently recommended in conventional canola production (1,000 mites/m<sup>2</sup>) (4). So, yield losses from ryegrass competition and RLEM damage were not expected to be significant. Despite the differences in canola emergence, there was no difference between the seeding rates in crop biomass production (Table 2) suggesting that the canola was able to compensate in biomass growth from initial low plant densities. Our results suggest that a higher canola seeding rate may provide sufficient competition to reduce weed biomass production, and thus have a positive effect on grain yield. Greater precision in seeding rate is required if these results are to be validated further in this environment.

**Table 1. Canola and ryegrass density at emergence (18/7/01) and RLEM density (15/6/01).**

Seeding rate (kg/ha)	Canola emergence (plants/m <sup>2</sup> )	Ryegrass density at emergence (plants/m <sup>2</sup> )	RLEM density (No./m <sup>2</sup> )
5	73	85	54
7.5	68	79	66
10	97	109	109
LSD P=0.05	19	ns	ns

**Table 2. Ryegrass density at early flowering of canola, canola and weed biomass, grain yield, oil content and grain protein.**

Seeding rate (kg/ha)	Ryegrass density at early flowering (plants/m <sup>2</sup> )	Canola biomass (kg/ha)	Weed biomass (kg/ha)	Grain yield (t/ha)	Oil content (%)	Grain protein (%)
5	107	2645	690	1.4	39.6	21.7
7.5	111	2774	860	1.3	39.2	21.9

<b>10</b>	106	2607	514	1.6	38.5	22.0
<b>LSD P=0.05</b>	<b>ns</b>	<b>ns</b>	<b>269</b>	<b>0.2</b>	<b>ns</b>	<b>ns</b>

## Conclusion

Higher seeding rates can result in higher canola yields in organic production systems, but reduction of RLEM population in the previous year and prior weed management is critical to achieve this result.

## Acknowledgments

The authors gratefully acknowledge the funding provided for this work by the Department of Natural Resources and Environment in Victoria under the Naturally Victorian Initiative.

## References

- (1) McCoy, S. and Parlevliet, G. (1998) The Export Market Potential for Clean and Organic Agricultural Products. A Report for the Rural Industries Research and Development Corporation, Agriculture Western Australia, Perth.
- (2) Peltzer, S. (1999) Controlling weed seed production with crop seeding rate. Weed Updates, Western Australia, 17-18 Feb 1999, Rendezvous Observation City WA.
- (3) Lemerle, D., Verbeek, B. and Coombes, N. (1995) Weed Research, 35, 503-509.
- (4) Anon. (1999) How many is too many? Insect Control Thresholds Grains Industry Advice, GRDC publication PP 504655/00130.